

Development of chemiluminescent methods for explosives detection

Elisabetta MAIOLINI^a, Stefano GIROTTI*^a, Elida N. FERRI^a, Pasquale CAPUTO^a, Gianluca GUARNIERI^b, Sergei A. EREMIN^c, Angel MONTOYA^d, Maria J. MORENO^d, and Marcello D'ELIA^e.

^a*Dipartimento di Scienza dei Metalli, Elettrochimica e Tecniche Chimiche, Università di Bologna, via San Donato 15, 40127 Bologna, Italy*

^b*Labo-Scientifica s.r.l., Parma, Italy*

^c*Department of Chemical Enzymology, Faculty of Chemistry, M.V. Lomonosov Moscow State University, Moscow 119899, Russia*

^d*Instituto Interuniversitario de Investigación en Bioingeniería y Tecnología Orientada al Ser Humano Universidad Politécnica de Valencia, Camino de Vera, s/n 46022 Valencia Spain*

^e*Gabinetto Regionale di Polizia Scientifica per l'Emilia Romagna, Via Volto Santo 3, 40123 Bologna, Italy*

Abstract. The present work was finalised to develop chemiluminescent assays more sensitive and easy to use than the analytical methods currently in use to detect traces of TNT (2,4,6-trinitrotoluene), TATP (Triacetone triperoxide) and HMTD (Hexamethylene triperoxide diamine). For TNT detection we developed an indirect competitive ELISA with chemiluminescent end-point (CL-ELISA) and a Lateral Flow ImmunoAssay (LFIA) for on-site analysis, based both on monoclonal antibody commercially available and conjugates specifically synthesized. For the peroxide-based explosives TATP and HMTD we developed an indirect assay which measures, by luminol emission, the H₂O₂ released by these compounds. Both methods showed good sensitivity and reproducibility, with detection limit, LOD, and IC₅₀ values in the ng mL⁻¹ range, CV values lower than 10%. This indirect assay was finally optimized for using in a portable luminometer.

Keywords: explosives, TNT, TATP, HMTD, chemiluminescence, CL-ELISA, LFIA, portable luminometer.