

Isomeric carbanion disubstituted ylides synthesis

Neculai Doru MIRON^{a*}, Denisa Ileana NISTOR^a, Ilie BUCUR^a and Gheorghe SURPĂȚEANU^c

^a*Department of Chemistry Engineering, University Bacău, 157 Calea Mărășești, 600115, Romania*
^b*Organic and Environnement Synthesis Laborator, “Côte d'Opale du Littoral” University of Dunkerque,
220 avenue de l'Université – BP 552659379 Dunkerque, France*

Abstract This paper studies the synthesis of two isomers carbanion disubstitute ylides: 3-phenacyl benzotriazolium-1-iu-4'-chlorobenzoil, 2'', 4'', 6''-trinitrophenyl-methylide and 1-4'-chlorophenacyl benzotriazolium-3-iu benzoil, 2'', 4'', 6''-trinitrophenyl-methylide. The synthesis starts with: benzotriazolium and 4 chlorine substituted ω -bromine acetophenone. The intermediary products obtained are: 1-N-phenacyl benzotriazolium, 3-phenacyl 4-chlorophenacyl benzotriazolium-1-iu bromide and in situ the tautomere carbanion monosubstituted ylides. To observe those ylides, as well as the size of 3:1 rapport, the ^1H -RMN spectra was used.

Keywords: synthesis, RMN analysis, benzotriazolium salts, isomers disubstituted ylides
